

On the volume of a six-dimensional polytope

A. Felikson, P. Tumarkin

Abstract. This note is a comment to the paper [7]. That paper concerns with the projective surface S in $\mathbb{P}^3$ defined by the equation $x_1x_2x_3 = x_4^3$. It is shown there that the evaluation of the leading term of the asymptotic formula for the number of rational points of bounded height in $S(\mathbb{Q})$ is equivalent to the evaluation of the volume of some 6-dimensional polytope $\mathcal{P}$. The volume of $\mathcal{P}$ is known from several papers; we calculate this volume by elementary method using symmetry of the polytope $\mathcal{P}$. We also discuss a combinatorial structure of $\mathcal{P}$.

1 Introduction

Consider the projective surface S in $\mathbb{P}^3$ defined by the equation

$$x_1x_2x_3 = x_4^3.$$

A few years ago several authors, [2], [4], [5], [7], [8], obtained an asymptotic formula for the number of rational points of bounded height in $S(\mathbb{Q})$, as a simple illustration of a rather general conjecture discussed in [1], [2], [8]. To explicitly evaluate the leading term of the asymptotic formula obtained in [2], [7], [8] one must calculate the volume $Vol(\mathcal{P})$ of the polytope $\mathcal{P} \in \mathbb{R}^6$ determined by the inequalities

$$\begin{aligned} x_{12} + x_{13} + 2(x_{21} + x_{31}) &\leq 1, \\ x_{21} + x_{23} + 2(x_{12} + x_{32}) &\leq 1, \\ x_{31} + x_{32} + 2(x_{13} + x_{23}) &\leq 1, \\ x_{ij} &\geq 0, \end{aligned}$$

where

$$(x_{12}, x_{23}, x_{31}, x_{13}, x_{32}, x_{21})$$

are Cartesian coordinates in $\mathbb{R}^6$.

As it is shown in papers [2], [7] and [8],

$$Vol(\mathcal{P}) = \frac{1}{4 \cdot 6!} = \frac{1}{2880}. \quad (1)$$

The goal of this note is to prove formula (1) by a direct elementary method, using symmetry of the polytope $\mathcal{P}$. We dissect $\mathcal{P}$ into 6 congruent parts and decompose one of these parts into 6 simplices. Then we show that the volumes

of these simplices sum up to $\frac{1}{24}Vol(\Delta)$, where Δ is the standard six-dimensional simplex determined by the inequalities

$$x_{12} + x_{23} + x_{31} + x_{13} + x_{32} + x_{21} \leq 1, \quad x_{ij} \geq 0, \quad i \neq j, \quad 1 \leq i, j \leq 3.$$

We use the notation and terminology of [6]. In particular, given a subset X of $\mathbb{R}^n$, $n \in \mathbb{N}$, we write $conv X$ and ∂X for the convex hull and the boundary of X respectively.

Acknowledgment. We would like to thank B. Z. Moroz who brought the problem to our attention, initiated numerous stimulating discussions and convinced us to write down this note.

2 Volume of a rational polytope

Let $\mathbb{E}^n$ be the n -dimensional Euclidean space with Cartesian coordinates. Let $\mathbf{v}_0, \mathbf{v}_1, \dots, \mathbf{v}_k$ be rational points in $\mathbb{E}^n$, i.e. the coordinates $(v_{i1}, \dots, v_{in})$ of $\mathbf{v}_i$ are rational numbers for all $i = 0, \dots, k$. Suppose that the points $\mathbf{v}_0, \mathbf{v}_1, \dots, \mathbf{v}_k$ are in the convex position, that is no of these points belongs to the convex hull of the others. Suppose that $k \geq n$. Then the set $conv\{\mathbf{v}_0, \dots, \mathbf{v}_k\}$ is an n -dimensional polytope with non-zero volume $Vol(\mathbf{v}_0, \dots, \mathbf{v}_k)$. In this section we describe how to find $Vol(\mathbf{v}_1, \dots, \mathbf{v}_n)$.

Denote by $det(\mathbf{v}_1, \dots, \mathbf{v}_n)$ the determinant

$$det(\mathbf{v}_1, \dots, \mathbf{v}_n) = \begin{vmatrix} v_{11} & v_{12} & \dots & v_{1n} \\ v_{21} & v_{22} & \dots & v_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ v_{n1} & v_{n2} & \dots & v_{nn} \end{vmatrix}.$$

If $k = n$, the polytope $conv\{\mathbf{v}_0, \mathbf{v}_1, \dots, \mathbf{v}_k\}$ is a simplex (may be degenerate). Recall that

$$Vol(\mathbf{v}_0, \mathbf{v}_1, \dots, \mathbf{v}_n) = \frac{1}{n!} \cdot det(\mathbf{v}_1 - \mathbf{v}_0, \mathbf{v}_2 - \mathbf{v}_0, \dots, \mathbf{v}_n - \mathbf{v}_0)$$

(see for example [3], Chapter 9).

Definition. A *triangulation* of a polytope is a decomposition of this polytope into finite number of simplices $T_1, \dots, T_r$, such that if $T_i \cap T_j \neq \emptyset$, then $T_i \cap T_j$ is a face of both T_i and T_j .

The formula above reduces the problem to the question of triangulation of the polytope $conv\{\mathbf{v}_0, \mathbf{v}_1, \dots, \mathbf{v}_k\}$.

Lemmas 1 through 3 are either well-known or elementary.

Lemma 1. Let $\mathbf{v}_i$, $1 \leq i \leq n$ be n points in general position in $\mathbb{E}^n$ (i.e. no k of these points belong to $(k-2)$ -plane). Let H be the hyperplane spanned by

$\mathbf{v}_1, \dots, \mathbf{v}_n$. Let $\mathbf{a}$ and $\mathbf{b}$ be two points in $\mathbb{E}^n$. The points $\mathbf{a}$ and $\mathbf{b}$ belong to different open half-spaces with respect to H if and only if

$$\det(\mathbf{v}_1 - \mathbf{a}, \mathbf{v}_2 - \mathbf{a}, \dots, \mathbf{v}_n - \mathbf{a}) \cdot \det(\mathbf{v}_1 - \mathbf{b}, \mathbf{v}_2 - \mathbf{b}, \dots, \mathbf{v}_n - \mathbf{b}) < 0.$$

Lemma 2. Let

$$S := \{Q, Q_i \mid 1 \leq i \leq m\}$$

be a set of $m + 1$ polytopes satisfying the following conditions:

- a) $Q_i \subseteq Q$ for $1 \leq i \leq m$;
- b) each facet of Q_i is a facet of one the polytopes belonging to the set $Q \setminus \{Q_i\}$ for $1 \leq i \leq m$.

Then

$$Q = \bigcup_{i=1}^m Q_i$$

Definition. Let P be a polytope in $\mathbb{E}^n$. A point $\mathbf{y} \in P$ is *visual* from the point $\mathbf{x} \notin P$, if $[\mathbf{x}, \mathbf{y}] \cap P = \mathbf{y}$ (where $[\mathbf{x}, \mathbf{y}]$ is the segment $\{\lambda \mathbf{a} + (1 - \lambda) \mathbf{b} \mid 0 \leq \lambda \leq 1\}$). A facet f of P is *visual* from the point $\mathbf{x} \notin P$, if each point of f is visual from $\mathbf{x}$.

If P is convex any facet containing at least one inner visual point is visual.

Lemma 3. Let P be a convex polytope, and $\mathbf{x} \notin P$ be a point. Any point $\mathbf{y} \in P$ visual from $\mathbf{x}$ is contained in at least one visual facet of P .

Theorem 1. Let $T = \bigcup_{i=1}^m T_i$ be a triangulation of $\text{conv}\{\mathbf{v}_1, \dots, \mathbf{v}_k\}$. Let $\mathbf{v}_0 \notin \text{conv}\{\mathbf{v}_1, \dots, \mathbf{v}_k\}$. Let $f_1, \dots, f_r$ be all facets of $T_1, \dots, T_m$ visual from $\mathbf{v}_0$. Denote $T_{m+j} := \text{conv } \mathbf{v}_0 \cup f_j$, $1 \leq j \leq r$.

Then $T' = \bigcup_{i=1}^{m+r} T_i$ is a triangulation of $\text{conv}(\mathbf{v}_0, \mathbf{v}_1, \dots, \mathbf{v}_k)$.

The proof is obvious.

Theorem 1 suggests an **algorithm** of triangulation of $\text{conv}\{\mathbf{v}_0, \mathbf{v}_1, \dots, \mathbf{v}_k\}$: Consider any $n + 1$ non-coplanar points (without loss of generality we may assume that these points are $\mathbf{v}_0, \mathbf{v}_1, \dots, \mathbf{v}_n$). To check that these points are non-coplanar make sure that $\det(\mathbf{v}_1 - \mathbf{v}_0, \dots, \mathbf{v}_n - \mathbf{v}_0) \neq 0$. Then $T_1 = \text{conv}\{\mathbf{v}_0, \mathbf{v}_1, \dots, \mathbf{v}_n\}$ is a simplex and T_1 is triangulated by itself. Use Theorem 1 to successively triangulate $\text{conv}\{\mathbf{v}_0, \mathbf{v}_1, \dots, \mathbf{v}_i\}$ for $i = n + 1, n + 2, \dots, k$.

3 Combinatorial structure of the polytope $\mathcal{P}$

Denote by α_1, α_2 and α_3 the hyperplanes

$$x_{12} + x_{13} + 2(x_{21} + x_{31}) = 1,$$

$$x_{21} + x_{23} + 2(x_{12} + x_{32}) = 1,$$

$$x_{31} + x_{32} + 2(x_{13} + x_{23}) = 1$$

respectively. Denote by π_{ij} the hyperplane $x_{ij} = 0$ ($i, j = 1, 2, 3, i \neq j$). Denote by σ the hyperplane $x_{12} + x_{23} + x_{31} + x_{13} + x_{32} + x_{21} = 1$.

It is easy to see, that if $X \in \sigma$ and $X \in \mathcal{P}$ then $X \in \alpha_i$ for $i = 1, 2, 3$. Therefore, σ is not a facet of $\mathcal{P}$. Thus, $\mathcal{P}$ is bounded by α_i , $i = 1, 2, 3$ and six facets π_{ij} . Any vertex of $\mathcal{P}$ is an intersection of at least 6 facets. Thus, any vertex of $\mathcal{P}$ is a subject to at least 6 equations under consideration. A straightforward calculation shows that $\mathcal{P}$ is a convex hull of 21 points. We list these points in Table 3. The right column of the table shows, to which of the hyperplanes α_j , $1 \leq j \leq 3$, if any, the corresponding point belongs. Note, that the indices indicate non-zero coordinates.

We present below the *Gale diagram* of $\mathcal{P}$, describing the combinatorial type of the polytope (see Figure 1).

$\mathcal{P}$ is a 6-dimensional polytope with 9 facets. The combinatorics of a convex n -polytope with $n + 3$ facets can be described by 2-dimensional *Gale diagram* (see [6]). This consists of $n + 3$ points $a_1, \dots, a_{n+3}$ of unit circle in $\mathbb{R}^2$ centered at the origin. The combinatorial type of a convex polytope can be read off from the Gale diagram in the following way. Each point a_i corresponds to the facet f_i of $\mathcal{P}$. For any subset J of the set of facets of $\mathcal{P}$ the intersection of facets $\{f_j | j \in J\}$ is a face of $\mathcal{P}$ if and only if the origin is contained in the set $\text{conv}\{a_j | j \notin J\}$.




Figure 1: Gale diagram of $\mathcal{P}$.

4 Triangulation of the polytope $\mathcal{P}$

In this section we use the group of symmetries of $\mathcal{P}$ to find a nice triangulation of $\mathcal{P}$ containing relatively small number of simplices.

Let

$$\mathcal{P}_j := \text{conv}\{O \cup \alpha_j\}, \quad 1 \leq j \leq 3.$$

Table 1: Vertices of $\mathcal{P}$.

Notation	Coordinates ($x_{12}, x_{23}, x_{31}, x_{13}, x_{32}, x_{21}$)	α_1	α_2	α_3
O	$(0, 0, 0, 0, 0, 0)$	—	—	—
Q_{23}^{21}	$(0, \frac{1}{2}, 0, 0, 0, \frac{1}{2})$	+	+	+
Q_{32}^{31}	$(0, 0, \frac{1}{2}, 0, \frac{1}{2}, 0)$	+	+	+
Q_{13}^{12}	$(\frac{1}{2}, 0, 0, \frac{1}{2}, 0, 0)$	+	+	+
P_{12}	$(\frac{1}{2}, 0, 0, 0, 0, 0)$	—	+	—
P_{23}	$(0, \frac{1}{2}, 0, 0, 0, 0)$	—	—	+
P_{31}	$(0, 0, \frac{1}{2}, 0, 0, 0)$	+	—	—
P_{13}	$(0, 0, 0, \frac{1}{2}, 0, 0)$	—	—	+
P_{32}	$(0, 0, 0, 0, \frac{1}{2}, 0)$	—	+	—
P_{21}	$(0, 0, 0, 0, 0, \frac{1}{2})$	+	—	—
R_1	$(\frac{1}{3}, \frac{1}{3}, \frac{1}{3}, 0, 0, 0)$	+	+	+
R_2	$(0, 0, 0, \frac{1}{3}, \frac{1}{3}, \frac{1}{3})$	+	+	+
T_{12}^{21}	$(\frac{1}{3}, 0, 0, 0, 0, \frac{1}{3})$	+	+	—
T_{23}^{32}	$(0, \frac{1}{3}, 0, 0, \frac{1}{3}, 0)$	—	+	+
T_{31}^{13}	$(0, 0, \frac{1}{3}, \frac{1}{3}, 0, 0)$	+	—	+
V_{21}^{32}	$(0, 0, 0, 0, \frac{1}{4}, \frac{1}{2})$	+	+	—
V_{12}^{31}	$(\frac{1}{2}, 0, \frac{1}{4}, 0, 0, 0)$	+	+	—
V_{13}^{21}	$(0, 0, 0, \frac{1}{2}, 0, \frac{1}{4})$	+	—	+
V_{31}^{23}	$(0, \frac{1}{4}, \frac{1}{2}, 0, 0, 0)$	+	—	+
V_{32}^{13}	$(0, 0, 0, \frac{1}{4}, \frac{1}{2}, 0)$	—	+	+
V_{23}^{12}	$(\frac{1}{4}, \frac{1}{2}, 0, 0, 0, 0)$	—	+	+

Lemma 4. $Vol(\mathcal{P}_1) = Vol(\mathcal{P}_2) = Vol(\mathcal{P}_3) = \frac{1}{3}Vol(\mathcal{P})$.

Proof. A permutation φ of coordinates

$$\varphi = (x_{12}x_{23}x_{31})(x_{13}x_{21}x_{32})$$

induces an order 3 orthogonal transformation $I_\varphi : \mathbb{R}^6 \rightarrow \mathbb{R}^6$, such that $I_\varphi(\mathcal{P}_1) = \mathcal{P}_2$, $I_\varphi(\mathcal{P}_2) = \mathcal{P}_3$ and $I_\varphi(\mathcal{P}_3) = \mathcal{P}_1$. Therefore,

$$\text{Vol}(\mathcal{P}_1) = \text{Vol}(\mathcal{P}_2) = \text{Vol}(\mathcal{P}_3). \quad (2)$$

By construction, the set $\{\mathcal{P}, \mathcal{P}_i \mid 1 \leq i \leq 3\}$ satisfies the conditions of Lemma 2, hence,

$$\mathcal{P}_1 \cup \mathcal{P}_2 \cup \mathcal{P}_3 = \mathcal{P}. \quad (3)$$

Moreover, if $1 \leq i < j \leq 3$, then $\mathcal{P}_i \cap \mathcal{P}_j$ is a common facet of $\mathcal{P}_i$ and $\mathcal{P}_j$; consequently,

$$\text{Vol}(\mathcal{P}_i \cap \mathcal{P}_j) = 0, \quad 1 \leq i < j \leq 3. \quad (4)$$

The lemma follows from (2)–(4). $\square$

In view of Lemma 4, it suffices to calculate the volume of $\mathcal{P}_1$. By construction, $\mathcal{P}_1$ is a convex hull of the points

$$\begin{aligned} &Q_{23}^{21}, T_{12}^{21}, P_{21}, R_1, V_{21}^{32}, V_{13}^{21}, \\ &Q_{32}^{31}, T_{13}^{31}, P_{31}, R_2, V_{31}^{23}, V_{12}^{31}, \\ &Q_{13}^{12}, O. \end{aligned}$$

Note that all vertices of $\mathcal{P}_1$ except for Q_{12}^{13} and O split into pairs of similar vertices denoted by one letter with different indices. This will be used in the proof of Lemma 5.

Denote by θ_1 and θ_2 respectively the hyperplanes

$$x_{12} - 2x_{23} + x_{31} = x_{13} + x_{32} - 2x_{21}$$

and

$$x_{12} + x_{23} - 2x_{31} = x_{13} - 2x_{32} + x_{21}.$$

It is easy to check that $\mathcal{P}_1$ has the following facets: $\pi_{ij}, \alpha_1, \theta_1$ and θ_2 .

Denote by δ the hyperplane

$$x_{12} + x_{23} + x_{31} = x_{13} + x_{32} + x_{21}.$$

Lemma 5. δ divides $\mathcal{P}_1$ into two congruent parts.

Proof. It is easy to check that $P_{31}P_{21}$ is the only edge of α_1 which is intersected by δ transversally.

Let $K_{21}^{31} := \delta \cap P_{31}P_{21} = (0, 0, \frac{1}{4}, 0, 0, \frac{1}{4})$. Then δ divides $\mathcal{P}_1$ into two polytopes $\mathcal{P}_1^1$ and $\mathcal{P}_1^2$, where $\mathcal{P}_1^1$ is a convex hull of

$$P_{31}, R_1, V_{31}^{23}, V_{12}^{31} \text{ and } O, Q_{23}^{21}, Q_{32}^{31}, Q_{13}^{12}, T_{12}^{21}, T_{13}^{31}, K_{21}^{31}$$

and $\mathcal{P}_1^2$ is a convex hull of

$$P_{21}, R_2, V_{21}^{32}, V_{13}^{21} \text{ and } O, Q_{23}^{21}, Q_{32}^{31}, Q_{13}^{12}, T_{12}^{21}, T_{13}^{31}, K_{21}^{31}.$$

Table 2: Triangulation of $\mathcal{P}_1^1$.

	Vertices	Neighbors	Volume
Δ_1	$O, P_{31}, Q_{32}^{31}, Q_{23}^{21}, T_{31}^{13}, T_{21}^{12}, K_{21}^{31}$	$\alpha_1, \delta, \pi_{32}, \pi_{23}, \pi_{13}, \pi_{12}, \Delta_2$	$\frac{1}{9} \cdot \frac{2}{4^3} \frac{1}{6!}$
Δ_2	$O, P_{31}, Q_{32}^{31}, Q_{23}^{21}, T_{31}^{13}, T_{21}^{12}, Q_{12}^{13}$	$\alpha_1, \delta, \pi_{32}, \pi_{23}, \Delta_3, \Delta_4, \Delta_1$	$\frac{1}{9} \cdot \frac{1}{4^2} \frac{1}{6!}$
Δ_3	$O, P_{31}, Q_{32}^{31}, Q_{23}^{21}, R_1, T_{21}^{12}, Q_{12}^{13}$	$\alpha_1, \theta_2, \pi_{32}, \Delta_5, \Delta_2, \Delta_4, \pi_{13}$	$\frac{1}{9} \cdot \frac{2}{4^2} \frac{1}{6!}$
Δ_4	$O, P_{31}, Q_{32}^{31}, Q_{23}^{21}, T_{31}^{13}, R_1, Q_{12}^{13}$	$\alpha_1, \theta_1, \pi_{32}, \pi_{21}, \Delta_3, \Delta_2, \Delta_6$	$\frac{1}{9} \cdot \frac{1}{4^2} \frac{1}{6!}$
Δ_5	$O, P_{31}, Q_{32}^{31}, V_{12}^{31}, R_1, T_{21}^{12}, Q_{12}^{13}$	$\alpha_1, \theta_2, \pi_{32}, \Delta_3, \pi_{23}, \pi_{21}, \pi_{13}$	$\frac{1}{9} \cdot \frac{1}{4^2} \frac{1}{6!}$
Δ_6	$O, P_{31}, Q_{32}^{31}, Q_{23}^{21}, T_{31}^{13}, R_1, V_{31}^{23}$	$\alpha_1, \theta_1, \pi_{32}, \pi_{21}, \pi_{13}, \pi_{12}, \Delta_4$	$\frac{1}{9} \cdot \frac{2}{4^3} \frac{1}{6!}$

Note that $\mathcal{P}_1^1$ and $\mathcal{P}_1^2$ differ by 4 vertices only.

Consider an involution I_ψ induced by the following permutation of coordinates: $\psi = (x_{12}x_{13})(x_{31}x_{21})(x_{32}x_{23})$. Clearly, I_ψ preserves α_1 and δ . More precisely, I_ψ fixes the points Q_{12}^{13} and O and interchanges six pairs of other points spanning $\mathcal{P}_1$. Thus, I_ψ interchanges $\mathcal{P}_1^1$ and $\mathcal{P}_1^2$. Since I_ψ is an isometry, $\mathcal{P}_1^1$ is congruent to $\mathcal{P}_1^2$. $\square$

Lemmas 4 and 5 show that the permutation group S_3 acts on $\mathcal{P}$. Moreover, S_3 is a group of symmetries of $\mathcal{P}$ and $\mathcal{P}_1^1 = \mathcal{P}/S_3$.

Lemma 5 implies that $Vol(\mathcal{P}_1^1) = Vol(\mathcal{P}_1^2)$. By Lemmas 4 and 5 it is sufficient to triangulate $\mathcal{P}_1^1$. This polytope is bounded by π_{ij} , α_1 , θ_1 , θ_2 and δ .

We decompose $\mathcal{P}_1^1$ into 6 simplices $\Delta_1, \dots, \Delta_6$ (see Table 4). We represent these simplices by their vertices. Each facet of Δ_i belongs either to some Δ_j ($i \neq j$) or to one of the hyperplanes bounding $\mathcal{P}_1^1$. Note, that a facet f of a simplex corresponds to the vertex $v(f)$ opposite to this facet and vice versa. For each vertex $v(f)$ of Δ_i we indicate the simplex Δ_j or the facet of $\mathcal{P}_1^1$ containing the facet f (see the third column).

Theorem 2. 1) $\mathcal{P}_1^1$ is triangulated by $\Delta_1, \dots, \Delta_6$.

2) $Vol(\mathcal{P}) = \frac{1}{4} \frac{1}{6!} = \frac{1}{4} Vol(\Delta)$.

Proof. The set vertices of $\Delta_1, \dots, \Delta_6$ coincides with the set of vertices of $\mathcal{P}_1^1$. Hence, the union of Δ_i ($i = 1, \dots, 6$) lies inside $\mathcal{P}_1^1$. The intersection $\Delta_i \cap \Delta_j$ is a face of Δ_i and Δ_j . It is left to show that any point of $\mathcal{P}_1^1$ belongs to some of Δ_i . The third column of Table 4 shows that each facet of $\Delta_i, i = 1, \dots, 6$ either belongs to $\partial\mathcal{P}_1^1$ or to some $\partial\Delta_j, j \neq i$. By Lemma 2, $\mathcal{P}_1^1 = \cup_{i=1}^6 \Delta_i$ and the first statement is proved.

By lemmas 4 and 5,

$$\text{Vol}(\mathcal{P}) = 6 \cdot \text{Vol}(\mathcal{P}_1^1) = 6 \sum_{i=1}^6 \text{Vol}(\Delta_i).$$

In view of triangulation shown in Table 4, this equals to $\frac{1}{4} \cdot \frac{1}{6!}$. This proves the first equality. The second equality is trivial. □

References

- [1] V. V. Batyrev and Yu. I. Manin, *Sur le nombre des points rationnelles de hauteur borné des variétés algébriques*. Math. Ann., 286 (1990), 27–43.
- [2] V. V. Batyrev and Yu. Tschinkel, *Tamagawa numbers of polarized algebraic varieties*. Astérisque, 251 (1998), 299–340.
- [3] M. Berger, *Geometry I*, Springer-Verlag Berlin Heidelberg, 1987.
- [4] R. de la Bretèche, *Sur le nombre des points de hauteur borné d’une certaine surface cubique singulière*. Astérisque, 251 (1998), 51–77.
- [5] E. Fouvry, *Sur la hauteur des points d’une certaine surface cubique singulière*. Astérisque, 251 (1998), 31–49.
- [6] B. Grünbaum, *Convex Polytopes*. Graduate Texts in Mathematics, Springer-Verlag, 2003.
- [7] D. R. Heath-Brown and B. Z. Moroz, *The density of rational points on the cubic surface $X_0^3 = X_1X_2X_3$* . Math. Proc. Camb. Phil. Soc. 125 (1999), 385–395.
- [8] P. Salberger, *Tamagawa measures on universal torsors and points of bounded height on Fano varieties*. Astérisque, 251 (1998), 91–258.

Independent University of Moscow, Russia
Max-Planck Institut für Mathematik Bonn, Germany
e-mail: felikson@mccme.ru pasha@mccme.ru